

AD-A051 923

CALIFORNIA UNIV SAN DIEGO LA JOLLA INST FOR PURE AND--ETC F/G 11/6  
A METALLURGICAL APPROACH TO HIGH TEMPERATURE SUPERCONDUCTIVITY.(U)  
FEB 78 B T MATTHIAS, Z FISK, D C JOHNSTON F49620-77-C-0009

UNCLASSIFIED

AFOSR-TR-78-0557

NL

|OF|  
AD  
A051923



END  
DATE  
FILMED  
5-78  
DDC

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS  
BEFORE COMPLETING FORM

1. REPORT NUMBER <b>AFOSR/TR-78-0557</b>	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A Metallurgical Approach to High Temperature Superconductivity.	5. TYPE OF REPORT & PERIOD COVERED FINAL 10/1/76-12/31/77	
6. AUTHOR(s) Bernd T./Matthias, Zachary/Fisk, David C. /Johnston <del>and</del> Robert N./Shelton		7. CONTRACT OR GRANT NUMBER(s) USAF/F49620-77-C-0009
8. PERFORMING ORGANIZATION NAME AND ADDRESS University of California, San Diego Institute for Pure and Applied Physical Sciences La Jolla, California 92093		9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102 F 16/2306/C1
10. CONTROLLING OFFICE NAME AND ADDRESS AF Office of Scientific Research (NE) Bolling AFB Bldg. 410 Washington, D. C. 20332		11. REPORT DATE Feb 1978
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Final rept. 1 Oct 76-31 Dec 77		13. NUMBER OF PAGES 30
14. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		15. SECURITY CLASS. (of this report) UNCLASSIFIED
16. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		17. DECLASSIFICATION/DOWNGRADING SCHEDULE
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Superconductivity Lattice Instability Metallurgy Electrical Resistivity		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A second system of ternary superconductors has been discovered. Their formula is $XRh_4B_4$ with X being almost any trivalent transition element. In addition the superconductor $ErRh_4B_4$ first becomes superconducting, then loses it and becomes ferromagnetic at a lower temperature. This phenomenon is a radically new feature for ordered compounds. The properties of these and some closely related superconductors and magnets have been investigated. We have further results on Chevrel-phase compounds, the first ternary superconducting system ever discovered: we find three distinct low temperature phases		

403 608.

LB

within the room temperature homogeneity range of  $\text{Cu}_x\text{Mo}_3\text{S}_4$ . A non-linear pressure variation of the superconducting transition temperature ( $T_c$ ) correlates with the crystallographic instabilities of  $\text{ZnMo}_5\text{S}_6$ ,  $\text{ZnMo}_5\text{Se}_6$  and  $\text{Cu}_{0.7}\text{Mo}_3\text{Se}_4$ .

A correlation between  $T_c$  and the composition induced metal-insulator transition in  $\text{AgSn}_{1-x}\text{Sb}_x\text{Se}_2$  has been found.

We provide an explanation for the low temperature resistance behavior of a number of high  $T_c$  A-15 superconductors, and a formalism for isolating the aspherical Coulomb scattering contribution to the electrical resistivity of metals has been developed. This leads to a principal understanding of the resistivity of superconductors above their transition temperature.

Low temperature x-ray measurements have been completed on crystallographically unstable systems  $(\text{La,Ce})\text{Ru}_2$  and  $(\text{Hf,Zr})\text{V}_2$ , and these instabilities correlated with  $T_c$ .

RECEIVED for	
RTIS	White Section <input checked="" type="checkbox"/>
SEC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
IDENTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

ACHIEVEMENT

Contract No: USAF/F49620-77-C-0009

Date: 10/1/76-12/31/77

A Metallurgical Approach to High Temperature Superconductivity

Bernd T. Matthias

University of California, San Diego

Research achievements for the period 1 October 1976 - 31 December 1977 are detailed below with reference to the "Statement of Work" of the proposal.

a) Work on New Multicomponent Superconductors

We have now discovered the second system of superconducting ternary compounds. They are the double Rh borides. The system is  $\text{XRh}_4\text{B}_4$  in which  $X = Y, \text{Nd}, \text{Sm}, \text{Er}, \text{Tm}, \text{Lu}$  and  $\text{Th}$ .<sup>1,2</sup> The superconducting transition temperatures ( $T_c$ ) are shown in Table I. For  $X = \text{Gd}, \text{Tb}, \text{Dy}$  and  $\text{Ho}$ , the isomorphous compounds become ferromagnetic instead (see Table I). Of particular interest are the high  $T_c$ 's of the  $\text{Er}$  and  $\text{Tm}$  compounds. Never before have such high superconducting transition temperatures been observed in compounds with elements having magnetic moments as high as  $9.6\mu_B$ .

The erbium compound is particularly interesting. The superconducting  $T_c$  is at 8.6K as pointed out before. Below 0.9K the compound loses its superconductivity and becomes magnetic instead.<sup>3</sup>

It is astonishing that even though the effective moments of  $\text{Ho}$  ( $10.6\mu_B$ ) and of neighboring  $\text{Er}$  ( $9.6\mu_B$ ) differ by only 10%,  $\text{HoRh}_4\text{B}_4$  is ferromagnetic whereas  $\text{ErRh}_4\text{B}_4$  is superconducting. Therefore, we explored the superconducting properties of the pseudoternary alloys  $(\text{Er}, \text{Ho})\text{Rh}_4\text{B}_4$  to clarify the reasons for this striking juxtaposition. Our results for this system<sup>4</sup>



are shown in Fig. 1. Here, we found that for up to 89% dilution of Er by Ho the superconducting transition temperatures are only weakly dependent on the Ho content and that the disappearance of superconductivity observed for pure  $\text{ErRh}_4\text{B}_4$  also persists over this concentration range. For higher Ho concentrations, only ferromagnetic transitions were observed. Other ternary systems in which superconductivity has recently been discovered include the Y-Os-Si<sup>5</sup> and Sc-Rh-Si<sup>6</sup> systems, with superconducting transition temperatures up to 10K and 9K, respectively. The composition and structure of each of the superconducting phases remain to be determined.

b) Work on Metastable Superconductors

Our work on the metallurgy of the above  $\text{RERh}_4\text{B}_4$  compounds led to the discovery of a structurally distinct high temperature modification (with the same stoichiometry) which could be obtained as a single phase by substituting small amounts of Ru for Rh. The superconducting and magnetic transition temperatures obtained for this body-centered-tetragonal series  $\text{RE}(\text{Rh}_{0.85}\text{Ru}_{0.15})_4\text{B}_4$  and for the pure Ru series of isostructural compounds,  $\text{RERu}_4\text{B}_4$ , are shown in Figs. 2a and 2b, respectively.<sup>7</sup> The  $T_c$ 's among the former series of compounds range from 4.0K for the Dy member to 9.6K for the Y compound, and no transitions into a magnetically ordered state were found above 1.5K; these materials thus constitute only the third known true ternary structure type to exhibit superconductivity in the presence of a large concentration of highly magnetic rare earth ions. Again, the presence of large concentrations of these highly magnetic rare earth ions does not severely interfere with the superconducting behavior.

A factor has been identified which correlates with the divergence in the superconducting properties of the two series of compounds shown in Fig. 2. Whereas the members of the superconducting  $\text{RE}(\text{Rh}_{0.85}\text{Ru}_{0.15})_4\text{B}_4$  series of compounds all exhibit  $c/a$  ratios less than 2.00, the  $\text{RERu}_4\text{B}_4$  series shows  $c/a$  ratios greater than 2.00, even though the unit cell volumes of corresponding members of each series are nearly the same (see Fig. 3). In order to study this correlation further and to determine the manner in which the shift from high  $T_c$ 's to low  $T_c$ 's comes about, we studied the crystallographic and superconducting properties of the  $\text{Y}(\text{Rh}_{1-x}\text{Ru}_x)_4\text{B}_4$  pseudoternary system. Our results,<sup>7</sup> shown in Fig. 4, indicate that the transition is quite abrupt, and that the transition does indeed occur at the composition ( $x \approx 0.5$ ) at which the  $c/a$  ratio increases past 2.00 with increasing  $x$ . The reasons for this correlation have not yet been identified.

$T_c$  data for the series of pseudoternary NaCl-type compounds  $\text{AgSn}_{1-x}\text{Sb}_x\text{Se}$ , shown in Fig. 5,<sup>8</sup> support the contention<sup>9</sup> that there exists some as-yet-unknown mechanism which enhances  $T_c$  for systems exhibiting abrupt composition-induced metal-semiconductor transitions. Whereas a monotonic decrease in  $T_c$  with increasing  $x$  is predicted on the basis of carrier concentration considerations,<sup>10</sup>  $T_c$  increases initially, reaches a peak for  $x = 0.15$ , and finally decreases precipitously to  $< 1.5\text{K}$  as a metal-semiconductor transition sets in at  $x = 0.4$ . This behavior is reminiscent of that observed previously for the  $\text{Li}_{1+x}\text{Ti}_{2-x}\text{O}_4$  spinel system.<sup>9,11</sup>

In the course of a continuing study of the relationship between high-temperature superconductivity and low-temperature lattice instability, we observed a very unusual variation of  $T_c$  with composition  $x$  (Fig. 6a) for the Chevrel-phase compounds  $\text{Cu}_x\text{Mo}_3\text{S}_4$ .<sup>12</sup> Results of low-temperature x-ray and electrical resistivity measurements (Fig. 6b and 6c) revealed that

the three  $T_c$  ranges of Fig. 6a are those of three distinct low temperature phases which are stable within the room temperature homogeneity range of rhombohedral  $\text{Cu}_x\text{Mo}_3\text{S}_4$ . Pressure-induced discontinuities in  $T_c$  were also found (Fig. 7), corresponding to transformations between the three phases. From the data of Figs. 6 and 7, the pressure-composition phase diagram in Fig. 8 was constructed.

c) Electrical and Magnetic Measurements on Superconducting Materials at Normal and High Pressures

We have identified one factor contributing to the anomalous pressure dependencies of  $T_c$  in the ternary molybdenum chalcogenides through hydrostatic compressibility measurements up to 29.8kbar on eleven of these compounds.<sup>13</sup> Data for the ternary sulfides are shown in Fig. 9. Analysis of the data reveals a small bulk modulus ( $B_0$ ) for each compound, with values comparable to those found in simple s-p elements such as indium. In contrast, the magnitude of the first pressure derivative of  $B_0$  is nearly three times as large as the largest reported values for elements.

The proposed relationship between the occurrence of nonlinear pressure variation of  $T_c$  and crystallographic instability<sup>14</sup> has been further verified through measurements on the Chevrel-phase compounds  $\text{ZnMo}_5\text{S}_6$ ,  $\text{ZnMo}_5\text{Se}_6$  and  $\text{Cu}_{0.7}\text{Mo}_3\text{Se}_4$ .<sup>15</sup>

Powder neutron diffraction data<sup>16</sup> have confirmed the existence of long-range magnetic order in the compound  $\text{HoMo}_6\text{S}_8$ . For this material, the type of order is ferromagnetic (Fig.10), commensurate with the crystallographic unit cell and is coincident with the destruction of superconductivity.



Thus, the competitive interplay between the two cooperative phenomena of magnetism and superconductivity yields similar results for  $\text{HoMo}_6\text{S}_8$  and  $\text{ErRh}_4\text{B}_4$ .<sup>3</sup> In contrast, low temperature neutron diffraction data<sup>17</sup> for  $\text{ErMo}_6\text{Se}_8$  reveal the presence of magnetic Bragg peaks which are not compatible with a simple ferromagnetic structure and this ternary compound remains superconducting below the temperature at which this magnetic structure occurs. Thus, ternary superconducting systems continue to display a rich variety of properties not previously observed in binary compounds.

One interesting feature of the magnetic behavior of the  $(\text{RE})\text{Rh}_4\text{B}_4$  compounds is that the Gd member does not have the highest magnetic ordering temperature, contrary to what one expects for magnetic exchange via conduction electrons. Other, simpler, borides (eg.  $(\text{RE})\text{B}_6$  and  $(\text{RE})\text{B}_4$ ) also behave in this same anomalous way. We are investigating this behavior in these simpler borides and have grown single crystals of several  $(\text{RE})\text{B}_4$ 's for magnetic and electrical measurements. These measurements are not complete.

Electrical resistivity measurements have been made on a number of the  $(\text{RE})\text{Rh}_4\text{B}_4$  compounds. Figure (11) shows the data for  $\text{HoRh}_4\text{B}_4$ . An anomaly at the magnetic ordering temperature is evident, as well as the negative curvature at higher temperatures, characteristic of superconducting materials.

We have found that an unusual phonon-spectrum (e.g., non-Debye like) is reflected, in the case of A-15 superconductors, by resistivity behavior which varies with  $T^2$  at low temperature.<sup>18</sup> We have observed that the resistivity of  $\text{Zr}_2\text{Rh}$  ( $T_c = 11\text{K}$ ) also fits such a power law above  $T_c$ , suggesting a similar cause for the unusual properties<sup>19</sup> of  $\text{Zr}_2\text{Rh}$ .



Theoretical work by Fulde and co-workers<sup>20</sup> predicts that scattering of conduction electrons by the aspherical part of the 4f electron orbits of rare earth ions can be a new coupling mechanism for superconductivity. We have presented a method for determining the magnitude of this scattering mechanism from electrical resistivity measurements on materials containing rare earth ions, and have successfully applied this method to  $\text{PrB}_6$  for which the effect was found to be relatively large.<sup>21</sup>

d) Low Temperature X-ray Diffraction Measurements

The occurrence of low-temperature lattice distortion in  $\text{Cu}_x\text{Mo}_3\text{S}_4$  for  $x \approx 2$  was found to increase  $T_c$  above that of the room-temperature rhombohedral phase.<sup>12</sup> This behavior is just the opposite to that previously observed for most other materials; for the latter, distortions to a less symmetric structure invariably reduced  $T_c$ .

Other systems exhibiting lattice transformations on which we have now completed measurements are  $(\text{La,Ce})\text{Ru}_2$  (correlation of  $dT_c/dp$  with the cubic-tetragonal transformation)<sup>22</sup> and  $(\text{Hf,Zr})\text{V}_2$  (detailed investigation of lattice distortions and their correlation with  $T_c$ ).<sup>23</sup>

REFERENCES

1. B. T. Matthias, E. Corenzwit, J. M. Vandenberg and H. E. Barz, Proc. Natl. Acad. Sci., USA 74, 1334 (1977).
2. J. M. Vandenberg and B. T. Matthias, Proc. Natl. Acad. Sci., USA 74, 1336 (1977).
3. W. A. Fertig, D. C. Johnston, L. E. DeLong, R. W. McCallum, M. B. Maple and B. T. Matthias, Phys. Rev. Letters 38, 987 (1977).
4. D. C. Johnston, W. A. Fertig, M. B. Maple and B. T. Matthias, Solid State Communications (in press).
5. C. Segre and D. C. Johnston, (unpublished results).
6. H. Braun (unpublished results).
7. D. C. Johnston, Solid State Communications 24, 699 (1977).
8. H. C. Ku and D. C. Johnston, (unpublished results).
9. D. C. Johnston, Ph.D. Thesis, University of California, San Diego (1975).
10. S. Geller, in Chemical Bonds in Solids, N. N. Sirota, ed., Vol. I (Consultants Bureau, New York, 1972), p. 43.
11. D. C. Johnston, J. Low Temp. Phys. 25, 145 (1976).
12. D. C. Johnston, R. N. Shelton and J. J. Bugaj, Solid State Communications 21, 949 (1977).
13. A. W. Webb and R. N. Shelton, J. Phys. F: Metal Phys. (in press).
14. R. N. Shelton, in Superconductivity in d- and f-Band Metals, (Plenum Press, New York, 1977) ed. D. H. Douglass, p. 137.
15. A. C. Lawson and R. N. Shelton, Mat. Res. Bull. 12, 375 (1977).
16. J. W. Lynn, D. E. Moncton, W. Thomlinson, G. Shirane and R. N. Shelton, Solid State Communications (in press).
17. J. W. Lynn, D. E. Moncton, G. Shirane, W. Thomlinson, J. Eckert and R. N. Shelton, Solid State Communications (in press).
18. G. W. Webb, Z. Fisk, J. J. Engelhardt and S. Bader, Phys. Rev. 15B, 2624 (1977).
19. S. L. McCarthy, Ph.D. Thesis, University of California, San Diego (1970).
20. P. Fulde, L. L. Hirst and A. Luther, Z. Phys. 230, 155 (1970).
21. Z. Fisk and D. C. Johnston, Solid State Communications 22, 359 (1977).

22. R. N. Shelton, A. C. Lawson and K. Babershke, Solid State Communications 24, 465 (1977).
23. A. C. Lawson, Phys. Rev. (in press).

TABLE I

Superconducting and magnetic transition temperatures of ternary  $\text{XRh}_4\text{B}_4$  compounds with a primitive tetragonal structure

Compound	Superconducting Transition Temperature (K)	Magnetic Ordering Temperature (K)
$\text{YRh}_4\text{B}_4$	11.34-11.26	
$\text{NdRh}_4\text{B}_4$	5.36-5.26	
$\text{SmRh}_4\text{B}_4$	2.51-2.45	
$\text{GdRh}_4\text{B}_4$		5.62
$\text{TbRh}_4\text{B}_4$		7.08
$\text{DyRh}_4\text{B}_4$		12.03
$\text{HoRh}_4\text{B}_4$		6.56
$\text{ErRh}_4\text{B}_4$	8.55-8.49	
$\text{TmRh}_4\text{B}_4$	9.86-9.73	
$\text{LuRh}_4\text{B}_4$	11.76-11.54	
$\text{ThRh}_4\text{B}_4$	4.34-4.29	
$\text{Lu}_{0.75}\text{Th}_{0.25}\text{Rh}_4\text{B}_4$	11.93-11.3	
$\text{Sc}_{0.75}\text{Th}_{0.25}\text{Rh}_4\text{B}_4$	8.74-8.49	






Fig. 1. Low temperature phase diagram for  $(\text{Er}_{1-x}\text{Ho}_x)\text{Rh}_4\text{B}_4$  compounds possessing the primitive tetragonal  $\text{YRh}_4\text{B}_4$  structure.




Fig. 2 Superconducting (o) and magnetic (o) transition temperatures for the isostructural series of compounds  $RE(Rh_{0.85}Ru_{0.15})_4B_4$  (Fig. 2a) and  $RERu_4B_4$  (Fig. 2b) which crystallize in the body-centered-tetragonal  $LuRu_4B_4$  structure. The arrows indicate temperatures above which no transitions were observed by ac magnetic susceptibility measurements, and the symbol X indicates that the element below it does not form the phase.




Fig. 3. Crystallographic data for two series of body-centered tetragonal compounds.




Fig. 4. Superconducting and crystallographic properties of the pseudoternary system  $Y(Rh_{1-x}Ru_x)_4B_4$ .






Fig. 5. Composition dependence of the superconducting transition temperature for the NaCl-type system AgSn<sub>1-x</sub>Sb<sub>x</sub>Se<sub>2</sub>.




Fig. 6 Superconducting transition temperature (Fig. 6a), lattice transformation temperature (Fig. 6b), and residual resistance ratio (Fig. 6c) vs. composition for  $\text{Cu}_x \text{Mo}_3 \text{S}_4$ .




Fig. 7 Hydrostatic pressure dependence of the superconducting transition temperature for  $\text{Cu}_x\text{Mo}_3\text{S}_4$ .




Fig. 8. Pressure-composition phase diagram for  $\text{Cu}_x \text{Mo}_3 \text{S}_4$  derived from the  $T_c$  data in Figs. 6a and 7.






Fig. 9. Compressibility data for Chevrel-phase sulfides.




Fig. 10. Magnetic diffraction pattern for  $\text{HoMo}_6\text{S}_8$ . Data taken at 1.00 K (above the magnetic ordering temperature  $T_M$ ) have been subtracted from the data taken at 0.05 K. The new Bragg peaks which develop below  $T_M$  coincide with the positions of the  $\text{HoMo}_6\text{S}_8$  nuclear peaks, establishing that the transition is to a state of long range ferromagnetic order.




Fig. 11. Temperature dependence of the electrical resistance of  $\text{HoRh}_4\text{B}_4$ .

Scientific Impact

The most important aspect of the work completed under this contract is the discovery of the occurrence of high superconducting transition temperatures in the presence of large localized magnetic moments. This extremely unusual phenomenon gives hope for higher  $T_c$ 's as well as the expectation now under investigation, that critical field measurements for these superconductors may provide important clues for the achievement of higher critical fields, a development which would be of great technologic significance.



STATEMENT OF WORK (1 October 1976 - 31 Dec. 1977)

- a) Synthesize, measure, and analyze new multicomponent superconductors.
- b) Synthesize and measure new metastable superconductors, by somehow circumventing their instabilities.
- c) Perform electric and magnetic measurements on superconducting materials at normal and high pressures.
- d) Perform low temperature x-ray diffraction measurements on superconductors.

Contract: AFOSR/F49620-77-C-0009

Date: 10/1/76 - 12/31/77

PUBLICATION LIST

1. Low Temperature Specific Heat and Superconductivity of Alpha-Phase Au-Ga Alloys, R. F. Hoyt, A. C. Mota and C. A. Luengo, Physical Review B14, 441 (1976).
2. The Effect of High Pressure on Superconducting Ternary Molybdenum Chalcogenides, R. N. Shelton, Proceedings of the Conference on Superconductivity in d- and f-Band Metals, ed. David H. Douglass, Rochester, N. Y. (April-May, 1976), pp. 137.
3. Inelastic Neutron Scattering Studies of the Phonon Spectra of Chevrel Phase Superconductors, S. D. Bader, S. K. Sinha and R. N. Shelton, Proceedings of the Conference on Superconductivity in d- and f-Band Metals, ed. David H. Douglass, Rochester, N. Y. (April-May, 1976), pp. 209.
4. Saturation of the High Temperature Normal State Electrical Resistivity of Superconductors, Z. Fisk and G. W. Webb, Proceedings of the Conference on Superconductivity in d- and f-Band Metals, ed. David H. Douglass, Rochester, N. Y. (April-May, 1976) pp. 545.
5. Calorimetric Observation of a Phase Transition in the Superconducting State in  $\text{Gd}_{1.2}\text{Mo}_6\text{Se}_8$ , R. W. McCallum, D. C. Johnston, R. N. Shelton and M. B. Maple, Proceedings of the Conference on Superconductivity in d- and f-Band Metals, ed. David H. Douglass, Rochester, N. Y. (April-May, 1976) pp. 625.
6. Some Surprises in Superconductivity, Bernd T. Matthias, Proceedings of the Conference on Superconductivity in d- and f-Band Metals, ed. David H. Douglass, Rochester, N. Y. (April-May, 1976) pp. 635.
7. Dependence of the Superconducting Transition Temperature on Electron per Atom Ratio and Lattice Deformation in Noble Metal Alloys, A. C. Mota and R. F. Hoyt, Solid State Communications 20, 1025 (1976).
8. Superconductivity, Bernd T. Matthias, Critical Materials Problems in Energy Production, Academic Press (1976), Chap. 21, pp. 663-682.

9. Superconductivity of New Metastable Phase of Scandium-Chromium, J. M. Vandenberg, B. T. Matthias, E. Corenzwit, and H. Barz, Journal of Solid State Chemistry 18, 395 (1976).
10. Specific Heat of a New Metastable Phase of Scandium-Chromium, R. W. McCallum, D. C. Johnston, M. B. Maple and B. T. Matthias, Mat. Res. Bull. 11, 781 (1976); 11, 1354 (1976).
11. Symmetries of Superconducting Sulfides, Bernd T. Matthias, International Journal of Quantum Chemistry 10, 435 (1976).
12. Measurement of the Pressure Dependence of  $T_c$  for Superconducting Spinel Compounds, R. N. Shelton, D. C. Johnston and H. Adrian, Solid State Communications 20, 1077 (1976).
13. Low Temperature Heat Capacity of Small  $Nb_3Sn$  Polycrystals by ac Calorimetry, R. Viswanathan and D. C. Johnston, Journal of Low Temperature Physics 25, 1 (1976).
14. Phonon Spectra of Chevrel-Phase Molybdenum Chalcogenide Superconductors, S. K. Sinha, S. D. Bader, G. S. Kanpp, G. E. Ostrowski and R. N. Shelton, Proc. Conf. on Neutron Scattering, Vol. I, 136-144, ed. R. M. Moon (1976).
15. Superconductivity of Neutron Irradiated  $Mo_3Os$ , A. R. Sweedler, S. Moehlecke, R. H. Jones, R. Viswanathan and D. C. Johnston, Solid State Communications 21, 1007 (1977); 23, vii (1977).
16. Superconducting and Normal State Properties of  $Ag_{1-x}Sn_{1+x}Se_{2-y}$ , D. C. Johnston and H. Adrian, Journal of Physics and Chemistry of Solids 38, 355 (1977).
17. Magnetic Properties of  $RE_xMo_6Se_8$  Compounds Between 0.7 and 295K, D. C. Johnston and R. N. Shelton, J. Low Temp. Physics 26, 561 (1977).

18. Comment on "Influence of Composition on the Superconducting Transition Temperature of Alloys with the A-15 Structure," G. W. Webb and B. T. Matthias, Solid State Communications 21, 193 (1977).
19. Pressure Dependence of  $T_c$  for the Shear-Stress-Induced Tetragonal Phase of  $V_3Si$ , R. N. Shelton, D. C. Johnston and R. Viswanathan, Mat. Research Bull. 12, 133 (1977).
20. High Superconducting Transition Temperatures of New Rare Earth Ternary Borides, B. T. Matthias, E. Corenzwit, J. M. Vandenberg and H. E. Barz, Proceedings of the National Academy of Sciences 74, 1334 (1977).
21. Aspherical Coulomb Scattering of Conduction Electrons in  $PrB_6$ , Z. Fisk and D. C. Johnston, Solid State Communications 22, 359 (1977).
22. Superconductivity, Lattice Transformations, and Electronic Instabilities in  $Cu_xMo_3S_4$ , D. C. Johnston, R. N. Shelton and J. J. Bugaj, Solid State Communications 21, 949 (1977).
23. The Crystallography of New Ternary Borides, J. M. Vandenberg and B. T. Matthias, Proceedings of the National Academy of Sciences 74, 1336 (1977).
24. Structural Transformation in Some Chevrel Phase Compounds:  $ZnMo_5S_6$ ,  $ZnMo_5Se_6$  and  $Cu_{0.7}Mo_3Se_4$ , A. C. Lawson and R. N. Shelton, Mat. Research Bull. 12, 375 (1977).
25. Superconductivity of Ternary Borides, Bernd T. Matthias, International Journal of Quantum Chemistry 11, 647 (1977).
26. Destruction of Superconductivity at the Onset of Long-Range Magnetic Order in the Compound  $ErRh_4B_4$ , W. A. Fertig, D. C. Johnston, L. E. DeLong, R. W. McCallum, M. B. Maple and B. T. Matthias, Phys. Rev. Letters 38, 987 (1977).



27. The Clustering Hypothesis of Some High-Temperature Superconductors, J. M. Vandenberg and B. T. Matthias, Science 198, 194 (1977).
28. Superconducting Transition Temperature, Its Pressure Dependence and Structural Transformation in (La,Ce)Ru<sub>2</sub> Alloys, R. N. Shelton, A. C. Lawson and K. Baberschke, Solid State Communications 24, 465 (1977).
29. Mode Softening and High Superconducting Transition Temperatures in Some A-15 Compounds, G. S. Kanpp, S. D. Bader and Z. Fisk, Ferroelectrics 16, 263 (1977).
30. Lattice Instabilities in non-A-15 Superconductors. A. C. Lawson and R. N. Shelton, Ferroelectrics 16, 73 (1977).
31. Ferroelectricity: Why Did It Take So Long?, Bernd T. Matthias, Ferroelectrics 16, 21 (1977).
32. Superconductivity in a New Ternary Structure Class of Boride Compounds, D. C. Johnston, Solid State Communications 24, 699 (1977).
33. LaAg Under Hydrostatic Pressure: Superconductivity and Phase Transformation, J. S. Schilling S. Methfessel and R. N. Shelton, Solid State Communications 24, 659 (1977).
34. Compressibilities and Volume Dependence of T<sub>c</sub> for Eleven Chevrel-Phase Superconductors, A. W. Webb and R. N. Shelton, J. Phys. F: Metal Phys. 8, 1 (1978).
35. Superconductivity Versus Ferromagnetism, F. Acker, B. T. Matthias and L. Rinderer, J. Low Temp. Phys. 30, 133 (1978).
36. The History of Superconducting Transition Temperatures, B. T. Matthias, U. S. Bureau of Mines, Amarillo, Texas (in press).
37. Normal-State Properties of Some Superconducting Ternary Molybdenum Sulfides, R. Viswanathan and A. C. Lawson, Indo-Soviet Conference on Solid State Materials, Bangalore, India (in press).

38. Low Temperature Crystal Structures and Superconductivity of  $(\text{Hf}_{1-x}\text{Zr}_x)\text{V}_2$  Alloys, A. C. Lawson, Phys. Rev. (in press).
39. Catalysis Versus Superconductivity, Bernd T. Matthias, Proceedings of the Willard Libby Workshop (in press).
40. The Conference: One Man's View, Bernd T. Matthias, Proc. International Conference on High Pressure and Low Temperature Physics (in press).
41. Pressure Dependencies of the Superconducting and Magnetic Critical Temperatures of Ternary Rhodium Borides, R. N. Shelton and D. C. Johnston, Proceedings of the International Conference on High Pressure and Low Temperature Physics (Plenum: New York, 1977) (in press).
42. LaAg Under Hydrostatic Pressure: Superconductivity and Phase Transformation, J. S. Schilling, S. Methfessel and R. N. Shelton, Proceedings of the International Conference on High Pressure and Low Temperature Physics (Plenum: New York, 1977) (in press).
43. Re-entrant Superconductivity and Magnetic Ordering in the Pseudoternary System  $(\text{Er}_{1-x}\text{Ho}_x)\text{Rh}_4\text{B}_4$ , D. C. Johnston, W. A. Fertig, M. B. Maple, and B. T. Matthias, Solid State Communications (in press).
44. Unusual Superconducting Behavior of the Molybdenum-Techetium System, A. L. Giorgi and B. T. Matthias, Phys. Rev. (in press).
45. Tetragonal Ternary Borides: Superconductivity, Ferromagnetism and the Role of Scandium, B. T. Matthias, C. N. K. Patel, H. Barz, E. Corenzwit and J. M. Vandenberg, Phys. Rev. Letters (in press).
46. Superconductivity of Transition Metal Sulfides, Selenides and Phosphides with NaCl Structures, A. R. Moodenbaugh, D. C. Johnston, R. Viswanathan, R. N. Shelton, L. E. DeLong and W. A. Fertig, submitted to Journal of Low Temperature Physics.

47. Neutron Diffraction Study of Magnetic Order in the Ternary Superconductor  $\text{ErMo}_6\text{Se}_8$ , J. W. Lynn, D. E. Moncton, G. Shirane, W. Thomlinson, J. Eckert and R. N. Shelton, to be published in Solid State Communications.
48. Direct Observation of Long Range Ferromagnetic Order in the Re-entrant Superconductor  $\text{HoMo}_6\text{S}_8$ , J. W. Lynn, D. E. Moncton, W. Thomlinson, G. Shirane and R. N. Shelton, to be published in Solid State Communications.
49. Superconducting and Magnetic Properties of  $\text{ErRh}_4\text{B}_4$ , H. R. Ott, W. A. Fertig, D. C. Johnston, M. B. Maple and B. T. Matthias, to be submitted to Journal of Low Temperature Physics.

COUPLING

1. High Transition Temperature Superconducting Materials

- a. Bernd T. Matthias, University of California, San Diego, IPAPS
- b. Consultation with other government agencies
- c. Los Alamos Scientific Laboratories, Los Alamos, New Mexico--The research on improving the basic properties of present superconducting materials for use in superconducting cables continues with several groups of researchers at Los Alamos Scientific Laboratories. For the most part, this coupling is still available only in classified form.
- a. Bernd T. Matthias, University of California, San Diego, IPAPS
- b. Conference
- c. Annual Meeting of the Southeastern Section of the American Physical Society
- a. Bernd T. Matthias, University of California, San Diego, IPAPS
- b. Special Lecture Series
- c. "Science in the Public Interest Lecture Series" sponsored by the University of Colorado, Boulder, Colorado

2. New Multicomponent Superconductors

- a. Bernd T. Matthias, University of California, San Diego, IPAPS
- b. APS March Meeting
- c. Announcement of  $\text{ErRh}_4\text{B}_4$  superconductor which also exhibits ferromagnetism.
- a. Bernd T. Matthias, University of California, San Diego, IPAPS
- b. Lecture Series
- c. Concepts in Modern Engineering and Technology, State University of New York at Binghamton, New York
- a. Bernd T. Matthias, University of California, San Diego, IPAPS
- b. Distinguished Lecturer
- c. University of Houston, Houston, Texas



3. Other

- a. Zachary Fisk, University of California, San Diego, IPAPS
- b. Collaboration
- c. A. Arko, G. Crabtree, D. Karim, F. M. Mueller, L. R. Windmiller, J. B. Ketterson, Argonne National Laboratories, deHaas van Alphen effect and the Fermi surface of  $\text{LaB}_6$
- a. Robert N. Shelton, University of California, San Diego, IPAPS
- b. Collaboration
- c. S. D. Bader, F. Y. Fradin, G. S. Kanpp and S. K. Sinha, Argonne National Laboratories, phonon spectra of Chevrel-phase superconductors
- a. Robert N. Shelton, University of California, San Diego, IPAPS
- b. Collaboration
- c. R. A. Hein and A. W. Webb, National Research Laboratories, Washington, D. C., compressibilities of Chevrel-phase ternary superconductors
- a. Robert N. Shelton, University of California, San Diego, IPAPS
- b. Collaboration
- c. S. Foner and E. J. McNiff, Jr., MIT National Magnet Laboratory, upper critical magnetic fields on ternary molybdenum selenides
- a. Robert N. Shelton, University of California, San Diego, IPAPS
- b. Collaboration
- c. J. W. Lynn, University of Maryland and D. E. Moncton, Bell Laboratories, Observation of long range magnetic order in ternary molybdenum chalcogenides
- a. Robert N. Shelton, University of California, San Diego, IPAPS
- b. Collaboration
- c. T. F. Smith, Monash University, Australia, Thermal expansion measurements on ternary molybdenum sulfides and selenides.
- a. David C. Johnston, University of California, San Diego, IPAPS
- b. Collaboration
- c. B. G. Silbernagel, Exxon Research Laboratories, NMR studies of ternary borides.
- a. David C. Johnston, University of California, San Diego, IPAPS
- b. Collaboration
- c. D. Taylor, Los Alamos Scientific Laboratories, Mössbauer measurements on  $\text{ErRh}_4\text{B}_4$ .